FOOD WASTE PREVENTION & MANAGEMENT

A guideline for South Africa











Food Waste Prevention & Management - A guideline for South Africa

DEPARTMENT OF ENVIRONMENT, FORESTRY AND FISHERIES

Design and layout by

Department of Environment, Forestry and Fisheries

Chief Directorate Communications and Advocacy Strategic Planning

Private Bag X447, Pretoria 0001

South Africa

DEFF & CSIR 2019, Food waste prevention & management: guideline for South Africa. Edition 1, DEFF & CSIR, Pretoria

Acknowledgment of the project team:	Ms Sihle Matinise (CSIR)
Mr Rishal Sooklal (DEFF)	Mr Tsebo Mohapi (DEFF);
Mr Dumisani Buthelezi (DEFF)	Mr Katlego Mabatle (DEFF); and
Prof Suzan Oelofse (CSIR)	Mr Kabelo Phakoe (DEFF).

CONTENTS

1. Setting the context: Food waste from the farm to the consumer	1
2. What is Food Waste?	1
3. Why is food waste important?	1
4. What is the purpose of this guide?	2
5. A snapshot of food waste	3
6. How much food is wasted at each stage of the supply chain?	4
7. What are the drivers for food wastage on farm?	6
8. What can farmers do to reduce on-farm food losses and waste?	8
9. What are the drivers for food wastage during processing?	8
10. What can food processors do to reduce food losses and waste?	9
11. What are the drivers for food wastage during distribution?	
12. What can food distributers do to reduce food losses and waste?	11
13. What are the drivers for food wastage during consumption?	11
13.1 What are the drivers of food wastage in households?	11
13.2 What can households do to reduce food waste?	13
13.2.1 Pre-Shopping	14
13.2.2 Shopping	14
13.2.3 Storage of Food Products	17
13.2.4 Handling Leftovers	20
13.2.5 Understanding best before and use by dates	20
13.3 What are the drivers of food wastage in the hospitality sector?	21
13.4 What can the hospitality industry do to reduce food waste?	23
14. Surplus food	26
15. Summary	26
16. References	27

ACRONYMS AND DEFINITIONS

Acronyms		
CSIR	Council for Scientific and Industrial Research	
DEA	Department of Environmental Affairs	
GDP	Gross Domestic Product	
FSC	Food Supply Chain	
FIFO	First In First Out	
SDG	Sustainable Development Goals	
US United States		
Definitions		
Consumer food waste	edible food produced for human consumption, but ends up thrown away	
Food loss	food loss refers to the decrease in food quantity or quality, which makes it unfit for human consumption	

Food waste	food waste is any food, and inedible parts of food, removed from the food supply chain to be recovered or disposed (including composted, crops ploughed in/not harvested, anaerobic digestion, bio-energy production, co-gen- eration, incineration, disposed to sewer, landfill or discarded to sea) and include food losses
Food surplus	Oversupply of edible food

WHAT IS IN THE GUIDELINE?

The guideline includes information on the drivers and possible actions that can be taken to prevent and manage food waste throughout the food supply. Information was extracted from various sources including journals, technical reports and guidelines. Only information containing definitions and statistics is referenced in the text. Sources of information that were used but not cited are included in the reference list.

The term 'food waste' is used in the broadest sense to include pre-consumer food losses and post-consumer food waste.



1. SETTING THE CONTEXT: FOOD WASTE FROM THE FARM TO THE CON-SUMER

Globally, it is estimated that one third of edible food is lost or wasted each year (Gustavsson *et al.*, 2011). South Africa generates about 10.2 million tonnes of food waste throughout the supply chain annually (Nahman and De Lange, 2013). Food waste comes with food insecurity as well as economic and environmental impacts. Efforts to reduce food waste therefore provide an attractive opportunity to improve food security in South Africa as well as to reduce the economic and environmental impacts associated with it.

KEY MESSAGES:

- No person should waste food, but most people waste food although often unintentional.
- Food waste reduction has become a paramount consideration to achieve sustainability due to the growing burden of global food insecurity coupled with economic and environmental impacts.
- Food waste reduction is important for contributing towards achieving the sustainable development goals (SDGs). The South African government has signed up to SDG 12.3 to reduce food waste by 50% by 2030.
- Efforts to reduce food waste requires active involvement of all the stakeholders across the food supply chain.
- Industry led initiatives such as the proposed Voluntary agreement developed by the Consumer Goods Council of South Africa are very important to reduce food waste.

2. WHAT IS FOOD WASTE?

Food waste is "any food, and inedible parts of food, removed from the food supply chain to be recovered or disposed (including composted, crops ploughed in/not harvested, anaerobic digestion, bio-energy production, co-generation, incineration, disposed to sewer, landfill or discarded to sea)" (Ostergen *et al.*, 2014). Some food waste is avoidable, while others are unavoidable. The inedible parts (peels, bones, pips, etc) of food that ends up as waste cannot be avoided, but the edible parts that ends up as waste is avoidable and should therefore not be wasted.

Surplus food is food that is fit for human consumption but which is produced or prepared in excess. If surplus food is not eaten by humans while it is fit for consumption, it also adds to food waste.

3. WHY IS FOOD WASTE IMPORTANT?

It is estimated that global food production must increase by 70% by 2050 to meet the demand for food. Yet, globally, nearly one third of all food produced for human consumption is lost or wasted each year Gustavsson et al., (2011). Food waste occurs at all stages of the food supply chain (FSC) from initial agricultural production on farm, through packaging, storage and processing, distribution and retail, up to the final consumer at household level.




Figure 1: The food supply chain

Wasting food has a triple negative effect:

- It negatively impacts on the economy as all water, electricity, seeds, fertiliser and other inputs used to produce the food is wasted if the food goes to waste. Food wastage therefore also impacts on water security.
- 2. It contributes to **food insecurity** by increasing the cost of food as the cost of the wastage gets factored into the prices of food, making food unaffordable for poor people.
- 3. It **contributes to climate change** by increasing greenhouse gas emissions into the atmosphere. The decomposition of wasted food disposed of at landfill generates methane, a greenhouse gas that is more effective at trapping heat in the atmosphere than carbon dioxide.

To support sustainable development, the United Nations have set a target to halve the per capita global food waste at the retail and consumer levels, and reduce food losses along production and supply chains, including post-harvest losses, by 2030.

Where in the supply chain the food is most wasted, depends on the country in question. It is generally accepted that developed countries like the US and Europe waste more food at the consumer stage of the supply chain while developing countries in Asia and Africa generate more food waste at the early stages of the supply chain, on farm and during packaging and processing. This behaviour is linked to the fact that consumers in industrialised countries are typically urbanised and buy food from supermarkets where abundance is portrayed. Consumers in developing countries are less urbanised, less reliant on supermarkets, and more likely to be involved in agriculture and primary food production. With the increasing urbanisation, income growth associated with economic prosperity, and resulting changes in dietary patterns, household food waste is likely to increase in South Africa and other developing countries alike if it is not addressed as a priority.

Food waste reduction is a priority for South Africa to address issues of food security and climate change while supporting the implementation of the waste management hierarchy (reduce, reuse, and recycle) and the Chemicals and Waste Phakisa.

4. WHAT IS THE PURPOSE OF THIS GUIDE?

This guide aims to assist South Africa to decouple economic development and food wastage at the consumer level of the supply chain. The purpose of this guide is therefore twofold:

 To raise awareness of food wastage throughout the supply chain, but specifically at consumer level in order to address food wastage before it reaches the same levels as in the developed world; and 2. To help all the players in the food supply chain to identify pain points where food waste is likely to occur, and to provide some pointers as to what can be done to prevent avoidable food waste and minimise unavoidable food waste.

5. A SNAPSHOT OF FOOD WASTE

A comparison of global versus South African food waste is provided in Table 1.

Table 1: Comparing South African and Global food wastage

Indicator	Global	South Africa
	TO ST	
	1.2	
Edible portion of food waste	1.3 billion tonnes per	10 million tonnes per year
	year	

Consumer food Europe and North On average households generate 6 ka per housewaste America hold per week = 95-115 kg per person per year On average people in Johannesburg dispose Sub-Saharan Africa of 12kg of food waste and South/Southeast per person per year into Asia the municipal bin and in Ekurhuleni 8 kg per = 6-11 ka per person person per vear per year Cost R37.57 trillion (1USD R75 billion (De Lange and =14.45 ZAR) (Jain Nahman, 2015) et al., 2018). About 2.2% of National GDP. R14.45 trillion (1USD =14.45 ZAR) of this R71.4 billion is opportunity cost is incurred cost and R3.6 billion of from environmental this cost is for disposal, impacts includina and externalities (social pesticide exposure, and environmental costs) water use, greenhouse gas emissions, loss of biodiversity and soil erosion.

Climate Change	4.4GT CO ₂ Equivalents from food waste is equal to 13.6% of global emissions from fuel combustion.	About 0.04GT of CO ₂ Equivalents from food waste which is 7.6% of the total greenhouse gas emissions of South African in 2010.
	2.8-4.14 tonne of CO ₂ emission equiv- alents per tonne of food wasted.	
	Globally, food waste is the third largest greenhouse gas emitter after China and the US.	
Water	The blue water foot- print of global food waste was 250 km³ in 2007. This is higher than the national blue water footprint account of any	1.7 km ³ of water is extracted from ground- water and surface water bodies to produce the food that was wasted in South Africa in 2012
	country. This is equivalent to 100 million Olympic sized swimming pools	This is equivalent to 680 thousand Olympic sized swimming pools or nearly 1/3 of the capacity of the Gariep Dam.

6. HOW MUCH FOOD IS WASTED AT EACH STAGE OF THE SUPPLY CHAIN?

The food wastage occur throughout the supply chain and the wastage differ depending on the type of commodity as well as site specific circumstances. As a general rule of thumb, soft and leafy fruit and vegetables are more likely to be wasted than roots and tubers. which are sturdier and not so easily damaged during handling and transportation. The estimated waste percentage for each commodity aroup in each step of the food supply chain for sub-Saharan Africa is provided in Table 2 below. The wastage in the early stages of the supply chain is expected to decrease with increased levels of sophistication applied during harvesting, and post-harvest handling and storage. Wastage during distribution is influenced by external factors such a road conditions, accessibility and proximity to markets. Poor road conditions contribute to bruising while long distance travel subtracts from the shelf life of commodities. Export is one example where commodities can be on-route to the market for up to three weeks on a ship, before reaching the consumer. It is likely that the wastage in South Africa at pre-consumer level is lower than the rest of sub-Saharan Africa, but this still needs to be confirmed.

Commodity group	Agricultural pro- duction	Post harvest han- dling and storage	Processing and packaging	Distribution	Consumer	Total
Cereals	6.0%	8.0%	3.5%	2.0%	1.0%	20.5%
Roots and Tubers	14.0%	18.0%	15.0%	5.0%	2.0%	54.0%
Oil seeds & Pulses	12.0%	8.0%	8.0%	2.0%	1.0%	31.0%
Fruits and Vegetables	10.0%	9.0%	25.0%	17.0%	5.0%	66.0%
Meat	15.0%	0.7%	5.0%	7.0%	2.0%	29.7%
Fish and Seafood	5.7%	6.0%	9.0%	15.0%	2.0%	37.7%
Milk	6.0%	11.0%	0.1%	10.0%	0.1%	27.2%

Table 2: Estimated waste percentage per commodity group at each stage of the supply chain for sub-Saharan Africa

The estimated edible portion of food waste throughout the supply chain in South Africa is estimated to be 10.2 million tonnes per year broken down into 2.7 million tonnes during agricultural production, 2.4 million tonnes during post-harvest handling and storage, 2.6 million tonnes during processing and packaging, 2 million tonnes during distribution and 0.5 million tonnes at the consumption stage. This wastage include locally produced and imported food as illustrated in Figure 2.




Figure 2: Estimated food waste throughout the supply chain in South Africa (from data from Nahman and De Lange, 2013)

Reported postharvest handling and storage losses in South Africa is provided in Table 3.

Commodity type	% Loss	Source
Cereals	15-30	DAFF, 2015
Roots and Tubers	10-40	
Oilseeds and Pulses	15-30	
Fruit and Vegetables	15-44	
Meat	6-8	
Milk	8-16	

Table 3: Postharvest handling and storage losses in South Africa

7. WHAT ARE THE DRIVERS FOR FOOD WASTAGE ON FARM?

The food losses and wastage on farm are the result of many different factors, some within the control of the farmer, and others not. South African farmers are categorised into commercial farmers and subsistence farmers. The commercial farmers typically supply fresh produce into the market, while subsistence farmers produce food for own consumption. The food losses and wastage on small scale farms without access to sophisticated harvesting, post-harvest hanaling, and storage facilities are generally higher than on the large commercial farms. The main drivers for food losses and wastage on farm are summarised in Table 4. **Case study:** Loss of food due to lack of infrastructure at Tshakhuma fruit market in Limpopo (Mashau *et al.*, 2012)

- 43.3% of the fruits is lost during post-harvest.
- Reasons for this wastage included:
- lack of refrigeration (the market is exposed to temperatures of 35-40°C in summer and 25-34°C in winter), Knowledge gaps in the use of proper packaging materials, poor procurement planning and the lack of understanding of the link between transport and fruit rotting.
- Food was not lost during transportation due to short distance

Driver	Cause	Result	
Weather	DroughtFloods	Damage to crops	
	 Hail Wind Heat waves 	Stress to animals Reduced quality of the food	
	• Cold spells	Delayed harvest- ing if the fields are inaccessible	

Table 4: Drivers of on-farm food wastage

Driver	Cause	Result
Harvest	 Mechanical damage 	Damage to crops
	 Spillage during harvest Crops left behind due 	Reduced quality of
	to poor harvesting	food
	techniques	Crops ploughed
	Crops not harvested due to price drops	back in
Disease and pests	Animal deaths during breeding	Condemnation at slaughterhouse
	Animal sickness	
	Disease of crops	Milk discards
	Crops eaten or dam- aged by pests	Crops ploughed back in
		Reduced quality or discards
Demand forecasting	Uncoordinated pro- duction i.e. all farmers produce tomatoes	Oversupply at markets
Grading	Grading errors	Rejected at market
-	• Out-grades	
Storage	 Packaging failure 	Spillage, reduce
	 Interruption in cold chain 	quality, or discards

Driver	Cause	Result
Transport	• Bruising	Reduced quality
	 Package failures 	Shortened shelf life
	 Exposure to the ele- ments 	Discarded
	 Cold chain failures 	
	 Road accidents 	
By-catch	 Non-target species caught by fisheries 	Discarded or pro- cessed as animal feed

8. WHAT CAN FARMERS DO TO REDUCE ON-FARM FOOD LOSSES AND WASTE?

It is in the best interest of the farmer to reduce food losses and wastage as it impacts on their income potential. It is therefore important that the workers on the farm are also aware of the impact of their actions on the quality of the food produced.

A few general tips for farmers to consider include:

- Engage with potential markets to establish market needs and quality specifications.
- Diversify markets to ensure market access for lower quality produce in case of rejections by high end markets.
- Optimise harvesting times and techniques to preserve product quality and shelf life.
- Optimise post-harvest handling and storage based on the commodity type being produced.

- Optimise transport between farm and distribution. It should be noted that poor road conditions, even short stretches on which produce is transported, contributes to losses due to bruising.
- Imperfect or sub-standard commodities creates opportunities for small business development in agro-processing.

Extend the shelf life of the commodities by applying best practice during production

9. WHAT ARE THE DRIVERS FOR FOOD WASTAGE DURING PROCESS-ING?

Food processors receive the commodities from the farm or the market for processing. Fruit are typically canned or processed into juice. Vegetables, fish and meat are processed into frozen, canned or bottled products, or converted into convenience foods such a pre-prepared meals requiring minimum further preparation by consumers. The main drivers for food losses and waste in the processing part of the supply chain are summarised in Table 5.

Table 5: Drivers of processing food wastage

Driver	Cause	Result
Rejected input	 Product rejected for not meeting quality specifi- cation 	Discarded

Driver	Cause	Result
Losses in process	 Food safety concerns Inefficient process flows Accidental spillages Washing, peeling, slicing, and boiling 	Condemnation Discarded
	 Machine failures Trimming Process interruptions e.g. power outages Maintenance runs 	
Quality Specification	Over-demanding quality specifications	Discarded
Labelling errors	Incorrect information on printed labelSkewed labelling	Discarded
"Off-spec" pro- duction	• Poor product formu- lation	Discarded

10. WHAT CAN FOOD PROCESSORS DO TO REDUCE FOOD LOSSES AND WASTE?

The majority of the food wastage in the processing stage of the supply chain occur during the preparation stages. A significant amount of the waste, including the pips and peels, is believed to be unavoidable waste. Avoidable waste streams should be minimised by optimising process flows, stock rotation, and general good housekeeping. Trimmings and off-cuts should be used in secondary products where possible.

Some steps that processors can take to reduce food wastage include:

- Measure food losses and identify the causes
- Train staff to prevent food losses
- Process optimisation through:
 - Identifying hotspots and introduction of quick-win solutions to prevent wastage;
 - Putting measures in place to allow for rapid retrieval, reworking of products, or reintroduction of primary materials;
 - Finding solutions for production interruptions;
 - Effective stock rotation and controls;
 - o Coordinating production with clients and suppliers

11. WHAT ARE THE DRIVERS FOR FOOD WASTAGE DURING DISTRIBU-TION?

The distribution stage in the supply chain includes wholesale markets, supermarkets, retailers, and fish markets. In the South African situation it also includes informal vendors and spaza shops. The main drivers for food losses and waste in the distribution part of the supply chain are summarised in Table 6.

How much	food is	wasted in	this stage?
----------	---------	-----------	-------------

Case study: Vegetable losses at retail level in Stellenbosch (Munhuweyi, 2012)

- Revealed that 14.46% of tomatoes, 21.21% cabbage and 17.93% of carrots was lost in retails.
- Reasons for losses included mechanical injury and decay

Case study: Self-reported losses of food by vendors (Mashau et al., 2012)

- Vendors selling guava, avocado and pawpaw self-reported 50% of food waste in a study that was conducted in Tshakhuma fruit market, Limpopo.
- Losses occurred as a result of over-ripening due to a lack of measures to control the ripening processes of the fruits

Table 6: Drivers of distribution food wastage

Driver	Cause	Result
Distribution system	 Long travel distances Sub-optimal logistics Uneven road surfaces Road accidents 	Reduced shelf life Discard

Driver	Cause	Result
Date labels	 Food not sold in time Food items not used in time 	Donations Discarded
	 Poor stock rotation in-store/depot 	
Ordering systems	Over delivery	Donations
	Ordered too much	Discarded
Packaging	 Packaging failures 	Discarded
Product recalls	 Food safety concerns Compromised quality 	Condemnation
Quality specifications	Product rejected for	Donated
	being below quality	Discarded
Cold chain failures	 Fridges and freezers not operated at optimum temper- atures 	Condemnation
	 Power outages 	
	 Consumers moving stock out of cold chain areas 	

Driver	Cause	Result
Retail over-stocking	 Company policies requiring full shelves at all times 	Donations Discarded
Failure to distribute edible surplus	 Reputational concerns Strict liability Logistical challenges for redistribution 	Condemnation Discarded

12. WHAT CAN FOOD DISTRIBUTERS DO TO REDUCE FOOD LOSSES AND WASTE?

Food losses and waste at the distribution part of the supply chain are the highest value food losses due to the input required to get it to this stage in the supply chain. Reducing losses at this stage of the supply chain is likely to have a significant economic impact. A few general tips for wholesalers and retailers to consider include:

- Support local producers to keep transport distances for distribution to a minimum;
- Avoid uneven road surfaces where possible to reduce bruising of fresh fruit and vegetables;
- Optimise ordering systems, cold chain management, and stock rotation;
- Do away with overstocking, it is okay if the shelf is not always filled to the brim;
- Reduced price offering on stock nearing its sell by date to entice consumers to buy the food;

- In-store processing of produce that is past 'sell-by' but within 'use by' date into readymade salads, bakery items, prepared meals etc.; and
- Donate surplus food to organisations such as Food Forward South Africa.

13. WHAT ARE THE DRIVERS FOR FOOD WASTAGE DURING CONSUMP-TION?

The consumption stage of the supply chain includes consumption at household level, and in the hospitality sector where food is prepared and served out of home. The hospitality sector includes the commercial service sector (restaurants, fast food outlets, hotels and cafés in the formal sector) and the institutional service sector (universities, prisons, school and hospitals and work cafeterias).

13.1 What are the drivers of food wastage in households?

Household food waste is usually associated with multiple actions that consumers take when providing, handling, preparing and eating food. The actions increase or decrease the likelihood of food being wasted. Hence, food waste is often an unintended consequence of daily activities that are undertaken around food in a household. Drivers of food waste in the households are summarised in Table 7.

How much food is wasted at household level?

A Study based on municipal solid waste characterisation studies using bulk sampling with randomised grab sub-sampling conducted over a 6-week period during summer in 2014 (Johannesburg) and 2016 (Ekurhuleni) showed that households members dispose about 8 kg in Ekurhuleni and 12 kg in Johannesburg per person annually into the municipal bin (Oelofse *et al.*, 2018).

Household food wastage in the City of Tshwane was measured at an average of 6kg per household per week. The households separated the food waste for weekly collection and measurement by the project team (Ramukhwatho, 2016).

Table 7: Drivers of household food wastage

Cause	Result		
 Buying too much 	Expired food		
 Preparing too much 	Leftovers		
 Serving too much 	Lenoveis		
 Impulse buying 	Plate waste		
 Falling for special offers 			
 Poor stock control at home 			
 Food not in season 			
	 Buying too much Preparing too much Serving too much Impulse buying Falling for special offers Poor stock control at home 		

Driver	Cause	Result
Food preparation	 Food safety concerns Accidental spillages Washing, peeling, slicing, and boiling Trimming Cooking disasters (burnt, too much salt, wrong ingredients) Inability to use leftovers 	Discards
Fussy eating	 Allergies Picky eaters	Discards
Labelling	 Misinterpretation of un-pack labels Confusion – sell-by, use-by, best before, expiry 	Discards
Packaging	 Pack sizes not suited for household size Packaging does not empty easily Packet does not reseal 	Discards

Driver	Cause	Result
Storage	 Not knowing how to store fresh produce appropriately 	Short shelf life
	 Load shedding inter- rupting cold chain 	
Lifestyle influences	 Unplanned change in diet 	
	Cultural practices	

13.2 What can households do to reduce food waste?

Consumers are the end users of the food supply chain and therefore their preferences determine the demand, which in turn influence the supply of certain food types into the market. It can therefore be said that consumer demand is a huge driver of food wastage throughout the supply chain. In this regard, consumers are increasingly buying convenience foods, which require less input at home. This trend push preparation wastage to the earlier stages of the supply chain, while increasing the packaging waste at the consumer stage, but it also reduces the shelf life of the final product.

Households can reduce food waste by being vigilant during pre-shopping, shopping, storage, and preparation of food. A summary of household food waste prevention measures are provided in Table 8.



Table 8: Tips for households to consider

Food waste pre	evention and management measures of households
Planning	Check your refrigerator / cupboard to identify what you already have
	Make a shopping list (list should be done in conjunction with meal planning)
Shopping	• Stick to the shopping plan
	Avoid impulsive buying
	Buy from small local shops or grow your own food
	Buy seasonal food (Refer to Table 9)
	Buy small amounts of food enough for meals that day each time
	Avoid buying in bulk
Storage	Store dry food in a cool and dry place
	• All perishable food must be refrigerated except for potatoes, onion and banana.
	Cheese must be stored in a cupboard and refrigerated once opened
	Store food under correct temperature
	• recommended temperature for storing food is 0°C -4.4° C for refrigeration and -17.7°C for freezing
	Practice stock rotation in both cupboard and refrigerator
	Always keep your storage places organized
Preparation	• Plan meals
	Read recipes on how to prepare food
	Avoid preparing excessive food
Leftover	Store in an air tight container and refrigerate or freeze
	• Do not keep food at room temperatures for more than two hours after you have prepared the food
	Keep leftover food for 3-4 days
	• Find recipes to use your leftovers as ingredients of a new meal

	Once options for prevention are exhausted, manage food waste
ė	Feed people
l waste	Feed animals
food	Compost
nage	Anaerobic
Wan	Landfill

13.2.1 Pre-Shopping

Planning before shopping focussing on needs, is one strategy that can be used to reduce food waste at consumption level. This involves checking the food that you already have in the cupboards or refrigerator and making a shopping list in order to avoid buying more than what you actually need. At the same time, checking inventories also helps in making sure that the products that you are going to buy will have space in the cupboards and refrigerators. Making a shopping list is important to reduce food waste, but only if you are going to stick to the list. Shopping lists should be made in conjunction with meal planning and communication between different household members to avoid more than one family member, buying the same products by on the same day.

13.2.2 Shopping

Factors contributing to food waste include: source of food, frequency of buying, not sticking to a list in the store and buying in bulk.

Sticking to the list is very important as it reduces overbuying and buying things that you already have. With regards to source of food, consumers that do shopping in large supermarkets are likely to generate a lot of food waste compared to consumers that shop in local small shops or those who grow their own food.

Buying locally produced and seasonal foods reduces the use of resources compared to imported and non-seasonal fruits and vegetables as they require transportation and energy. A guide that can be used pre-shopping to check seasonal fruits and vegetables is provided in Table 9.

Buying small amounts of food enough for meals that day each time reduces food waste at household level. Buying in bulk increases the likelihood of food being wasted as a result of not using it on time.

Deciduous fruit Oct Nov Dec Jan Feb Mar Apr Mav Jun Jul Aug Sept Х х х х х х х х Х х Х Х х Х х Х х х х х х х х х Х х Х Х Х Х х Х Х х Х Х Х Х Х Х Х х Х Х Х х х х х х х Citrus Fruit Sept Nov Dec Feb Mar Apr May Aug Х х х х х Х х Х Х х Х х х Х Х Х Х Х Х Х Х Х Х Х Х х Х Х Sub-tropical fruit Feb Mar Sept Dec Jan Apr May Aug х х Х х Х х Х х х х х Х Х Х

Table 9: A guide that can be used pre-shopping to check for seasonal fruit and vegetables

$\langle 1/\rangle$

Mangoes		х	х	х	х	Х						
Pineapples	х	х	х	х	х	х	х	х	х	х	х	
Passion fruit					х	x	х	х	х	х	х	
Exotic fruit	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept
Raspberries		х	х	х	х	х						
Blueberries		х	х	х	х	x						
Melons		х	х	х	х	x						
Strawberries	х	х	х									
Figs			х	x	х	x	х	x				
Pomegranates					х	х	х	х	х			
Vegetables	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept
Brinjal			х	х	х	x	х					
Parsnips	х						х	х	х	х	х	х
	х	х	х	х	х	х	х					
Kale								х	х	х	х	
Baby marrow	х						х	х	х	х	х	х
Spinach	х	х	х	х	х	х	х	x	х			
	х				х	х	х	х	х	х	х	х
Mushroom	х	х	х	х				х	х	х	х	х

Rhubarb	х	х	х									х
Spring onions										х	x	х
Cauliflower	х	х					х	х	х	х	x	х
Broccoli	х						х	х	х	х	x	х
Green beans	Х	х	х	х	х	х	х	х				х
Broad beans										х	х	х
Peas	Х	х								х	x	х
Mangetout	Х	х	х	x	х	х						
Leeks	Х	х				х	х	х	х	х	х	х
Artichoke	х	х	х	х	х							х
Asparagus	х	х	х	х	х						х	х

13.2.3 Storage of Food Products

Food storage plays an important role in reducing the amount of food waste. Food storing includes dry storage and cold storage. Dry storage refers to storing of food in cupboards or a pantry while cold storage refers to storing food in the refrigerator or freezer. Improper storage or storing food for a long time are major contributors of food waste.

Refrigeration and freezing plays a critical role in reducing food waste as it prolongs the shelf life of a product. This food waste reduction technique is more important for food products that perish quickly such as fruits and vegetables. All fruits and vegetables except for bananas, onion and potatoes, should be refrigerated for prolonged shelf life. Depending on the type, vegetables can be kept for up to two weeks when refrigerated and up to 12 months when frozen. Critically important to refrigeration and freezing of food is the temperature as it affects microbial growth and the maturation, ripening and aroma development of fruits and vegetables. The recommended temperature for storing food is 0°C -4.4° C for refrigeration and -17.7°C for freezing. It is therefore important to ensure that food is stored under correct temperatures.

It is also important to note that refrigeration or freezing does not stop food spoilage but it simply retards the growth of microorganisms and slows down chemical changes that affect quality. Table 10 provides tips and recommendations on how long food can be stored in your cupboards, refrigerators and freezers.

Table 10: Tips and recommendations on how to store food

Product	Cupboard or pantry	Refrigerator	Freezer	Comments
Bakery goods:	5-7 days	1-2 weeks	2-3 months	Bread should be refrigerat-
Bread				ed to extend the shelf life or frozen it if it is not going to
Waffles		4-5 days	1 month	be used quickly. The quality of frozen bread is much bet-
Cake	1-2 days	1 week		ter than the quality of bread stored in the refrigerator.
Vegetables:	2-3 weeks			Refrigeration and freezing
Sweet potatoes				is the best way to maintain quality and delay ripening
White potatoes	2-3 months			for fresh vegetables. It is recommended to blanch
Spinach		1-2 days	10-12 months	(partially cook) or cook
Tomatoes	Until ripe	2-3 days	2months	vegetables before freezing. Vegetables such as pota-
Mushrooms		2-3 days	10-12 months	toes, sweet potatoes must be kept at room tempera-
Carrots		3 Weeks	10-12 months	tures. Potatoes must not be
Cauliflower		3-5 days	10-12 months	washed before storing them as that can shorten their
Garlic	1 month	1-2 weeks	1 month	shelf life.

Product	Cupboard or pantry	Refrigerator	Freezer	Comments
Fruits:	1-2 days	3 weeks	(cooked)	Bananas will blacken when
Apples			8 months	refrigerated but the inside colour will not change.
Bananas	Until ripe	2 days	1 months	
			(whole, peeled)	
Strawberries		1-2 days	8 months	
Avocados	Until ripe	3-4 days		
Meat:		3-5 days	4-12 months	
Roasts, beef, veal, pork, lamb				
Sausage		1-2 days	1-2 months	
Fish and sea food:		3-4 days		
Smoked Fish			2 months	
Oysters, clams, mussels		1-2 days		
Cereal:	1-2 years			Cereals should be stored in
Pasta, dry				air tight containers
Rice, white	2 years			
Flour	6-12 months			

In both refrigerators and cupboards, stock rotation is very important. Use the 'First-In First- Out' (FIFO) storage method where new products are kept behind so that the old products are used first. Avoiding chaotic storage areas is also important in order to prevent the risk of forgetting food.

13.2.4 Preparation of food

Lack of food preparation skills is a key cause of consumer food waste. Providing education, training and skills on how to prepare different food products is important to reduce food waste. The best way to learn how to prepare food is to assist and observe when someone who knows how to prepare food is in action.

13.2.5 Handling Leftovers

Leftovers occur as a result of preparing more food than needed. Reasons for preparing more food may include poor portioning and cooking for later use. Food commodities that are likely to end up as leftovers are small amounts of food such as rice, potatoes and pasta. Leftover food can also be reduced by proper portioning of food during preparation.

Proper handling of leftover food is important to ensure food safety when it is stored for later consumption. Food should not be kept at room temperature for more than two hours after preparation and should be kept in the refrigerator for no longer than 3-4 days.

13.2.6 Understanding best before and use by dates

Alleviating confusion on date labels is critical for food waste preven-

tion. Food date labels are reportedly erroneously used by consumers as an indication of food safety and that results in loss of food that is still fit for consumption.

In South Africa, food date labelling is regulated by Foodstuff, Cosmetics and Disinfectants Act no. 54 of 1972 (as amended). According to the regulations food date labels should appear in a calendar format ("Day-Month-Year") with the minimum durability written as "best before" or BB and "use by" date written in full. There are various date labels that are used to mark different food products and these include:

- "Best before" date- indicating the date at which the product is at its optimum quality. According to the Foodstuff, Cosmetics and Disinfectants Act the products may be sold and consumed beyond this date.
- "Use by date- indicating the end of the estimated period for the product's best quality, after which the product may lose some of its quality attributes. The food product may not be sold after this date.
- "Date of manufacture"-indicating the day on which the food becomes a finished product.
- "Sell by date"- indicating the last day on which the product should appear on the retail store shelf and its aim is to inform the retailers about the date on which the product should be sold. Beyond this date, it is safe to consume the product.

It is important to note that date labels as contained in the Act is not an indication of food safety. Food safety is influenced by amongst other, the integrity of the packaging material as well as the effectiveness of the cold chain. Food that is within the "best before" date can for instance be unfit for consumption if the packaging failed or the cold chain was interrupted. To verify food safety, consumers are encouraged to inspect the product before use. The food is likely to be safe for consumption if all the following are in place:



- the container is intact and the seal not broken;
- the food content looks and smell still good for consumption no obvious signs of discoloration, texture change, mildew, bacteria etc.,
- the food taste as expected when taking a small bite.

13.3 What are the drivers of food wastage in the hospitality sector?

The hospitality industry generates a significant amounts of food waste, owing to the high prevalence of eating out among South Africans. This sector is made up of commercial food service sector and institutional service sector. The commercial food service sector includes hotels, all types of restaurants (restaurants, fast food restaurants and buffet type restaurants) and clubs while the institutional service sector includes transport services, health (public and private hospitals), educational institutions, and prisons. The main drivers for food losses and wastage in the hospitality sector are summarised in Table 11.

Driver	Cause	Result
Planning	 Inaccurate demand forecasting 	Over-preparation Donations
	 Use of margin of error Over purchasing of stock 	Pre-consumer food waste
	 Having many dishes on your menu 	Plate waste
	 Lack of menus for kids 	
	 New dishes on the menu 	
	• Time limits on how long the food should be stored after prepa- ration	

Table 11: Drivers of food wastage in the hospitality sector

ver	Cause	Result	Driver	Cause	Result
ood preparation	 Food safety concerns Use of improper equipment (e.g. some peelers produce thick peels) Accidental spillages Washing, peeling, slicing, and boiling Unnecessary trimmings 	Discards	Serving	 Displaying large amount of food in a buffet Accidental spillages A patient discharged in a hospital Large serving plates Children dishing up for themselves in a buffet 	Reduced she Donations Discards
	 Cooking disasters (burnt, too much salt, wrong ingredients) Inability to repurpose off-cuts Preparation of more food than the demand Large portion sizes Inefficiency by the staff members 		Storage	 Not knowing how to store fresh produce appropriately Load shedding inter- rupting cold chain Packaging failure Poor stock rotation Malfunction of the refrigerators 	Short shelf life Discards
abelling	 Confusion – sell-by, use-by, best before, expiry 	Discards			

Driver	Cause	Result
Eating	 Dissatisfaction with the taste of food Ordering an item that a consumers don't know 	Discards
	 Loss of appetite on patients in hospitals Picky eating especially children 	

sumption at home should be encouraged. It should however be noted that salads and vegetables are seldom included in the 'doggy bag'.

13.4 What can the hospitality industry do to reduce food waste?

Food waste reduction is very important in the hospitality sector for economic benefits and also to avoid loss of valuable resources. Although the hospitality industry is diverse in terms of the types of facilities and how food is provided, actions towards food waste reduction can be applied across all the facilities. Some of the measures that can be used by the hospitality sector to reduce food waste include accurate customer demand forecasting, training of staff members, careful menu planning, acknowledging staff members that reduce food waste and donating food that is not sold. Consumers should also be involved in reducing food waste because they also contribute to generation of post consumption waste (plate waste) (Figure 3). A summary of hospitality food waste prevention measures are provided in Table 12.

In the hospitality industry left overs occur in the consumption stages (plate waste) of the service. The South African practice of offering a 'doggy bag' to consumers to take their left over food with for con-





Figure 3: A meal served in a restaurant (left) and the plate waste after the meal (right)

Table 12: Tips for the hospitality sector to consider

Food	l waste prevention	and management measures in the hospitality sector
l waste	Planning	 Check your refrigerator / cupboard to identify what you already have Make a shopping list (list should be done in conjunction with meal planning) Accurate forecast demand Prepare food to meet the demand
Prevent food	Shopping	 Stick to the shopping plan Buy from local suppliers Buy seasonal food (see guide) Prepare food to meet the demand
	Receiving	• Ensure that all the food is in good condition upon arrival
Prevent food wast	Storage	 Store dry food in a cool and dry place All perishable food must be refrigerated except for potatoes, onion and banana. Cheese must be stored in a cupboard and refrigerated once opened Store food under correct temperature (see link) Recommended temperature for storing food is 0°C -4.4° C for refrigeration and -17.7°C for freezing Practice stock rotation in both cupboard and refrigerator Always keep your storage places organized
	Preparation	 Prepare food to meet the demand Follow standard operation procedure Portion food correctly Provide training to the Kitchen operators to ensure proper handling of food Acknowledge staff members that reduce food waste

Food	I waste prevention	and management measures in the hospitality sector			
	Serving waste	• The buffet style used in hotels should be replaced with a la carte serving style			
		• Buffet waste can be reduced through the use of small platters. A label to inform guests that they can return for more food must also be used			
		• Use tasting plates with small portions to allow people to taste the food before dishing larger portions reduce			
		• Educate university students about the impacts of their irresponsible food waste behaviours (forgetting to unbook meals) to avoid surplus food			
	Plate waste	Use of small plates to reduce buffet plate waste			
	Reduce portion sizes				
	Once options f	or prevention are exhausted, manage food waste responsibly by separating the food waste at source and following the food waste hierarchy			
	Feed people	Donate surplus food (food prepared but not served) to charities			
	Feed animal	Donate to a pig farm or feed pets			
		• Meat must not be fed to pigs to avoid the risk of transmitting viruses to pigs, but it can be processed into commercial animal feed			
aste	Compost	Compost fruit and vegetable scraps and coffee grounds			
food waste	Anaerobic digestion	Separate collection of waste for anaerobic digestion			
Manage	Landfill	• Throw in the bin only if no other option is available (last resort).			
2					



14. SURPLUS FOOD

According to Papargyropoulou et al., (2014) surplus food refers to food produced beyond our nutritional needs. In the consumption stages of the FSC, the terms surplus food and leftover are often used interchangeably and thus making it difficult to distinguish between the two terms. For the purpose of this guideline leftover food refers to food left on the plate including both untouched and unfinished food. Surplus food on the other hand refers to food that has been prepared but not served to the people. Surplus food in the hospitality industry occurs in the serving stages of the service due to poor demand forecast and consequent preparation of excessive food and use of margin of error. In the households surplus food may arise as a result of stockpiling food and over-preparation of food. Papargyropoulou et al., (2014) recommends supply of food that is needed to prevent surplus food and donation of food to the needy where it is not possible to prevent it.

In the hospitality industry it is very challenging to recover surplus food because of the short shelf life (i.e. 24h on average). The need to transport food from the food service to the receivers every day acts as a barrier towards recovering food and that results to extremely high logistics costs. Health regulations also negatively impact on the food service sector's ability to donate food due to the potential risks to human health.

Food Forward distributed 4 350 tonnes of surplus food to 600 registered beneficiary organisations in 2017 in South Africa. An estimated 14.5 million meals were served using surplus food from these donations. This surplus food was recovered from retailers, food manufacturers and farmers. Rivett-Carnac and Von Bornmann (2018) estimate that 40 000 tonnes of surplus food exist across all grocery retail stores in South Africa every year. Woolworths is one of the retail stores that distributes and donates surplus food to 1000 charities on an on-going basis annually. This includes food products that have gone past the "Sell By" date but before "Best Before" date (Woolworths, 2016). The surplus that is not recovered for redistribution is most often disposed of on landfills as food waste.

15. SUMMARY

Food waste is a huge challenge that threatens our food system as well as our economic and natural resources. This is particularly true for food wasted in the consumption stages of the food supply chain due to the cumulative amount of resources used to produce the final product. There is a great opportunity to prevent and manage food waste throughout the food supply chain but specifically in the consumption stages of the FSC through behaviour change and education and awareness.

Measures taken to prevent and manage food waste may include planning before shopping, sticking to the plan, understanding date labels and packaging, and growing one's own food. When options for prevention are exhausted, unavoidable food waste should be managed by feeding surplus food to people, and animals through processed animal feed. Other waste management options include nutrient upcycling, composting, anaerobic digestion and value extraction through alternative waste treatment solutions.

Industry led initiatives which are in line with the food waste guideline must be adopted and supported. The Voluntary agreements being developed by the Consumer Goods Council of South Africa is one such initiative.

16. REFERENCES

- Cronjé, N., Van der Merwe, I. and Müller, I.M., 2018. Household food waste: A case study in Kimberley, South Africa. Journal of Consumer Sciences, 46.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. A profile of the South African Cabbage market value chain. Department of Agriculture, Forestry and Fisheries, Pretoria.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. A profile of the South African Carrot market value chain. Department of Agriculture, Forestry and Fisheries, Pretoria.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. A profile of the South African Dry bean market value chain. Department of Agriculture, Forestry and Fisheries, Pretoria.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. A profile of the South African Onion market value chain. Department of Agriculture, Forestry and Fisheries, Pretoria.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. A profile of the South African Sweet Potato market value chain. Department of Agriculture, Forestry and Fisheries, Pretoria.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. A profile of the South African Potato market value chain. Department of Agriculture, Forestry and Fisheries, Pretoria.
- DAFF (Department of Agriculture, Forestry and Fisheries), 2015. Briefing note on the Agro-processing sector in South Africa.

Available at: http://pmg-assets.s3-website-eu-west-1.amazonaws. com/151027brief.pdf

- Gustavsson, J., Cederberg, C., Sonesson, U., van Otterdijk, R., 2011. Global Food Losses and Food Waste; Extent, Causes and Prevention. Swedish Institute for Food and Biotechnology (SIK), Gothenburg (Sweden), and FAO, Rome (Italy).
- Marais ML, Smit Y, Koen N and Lötze E. 2017. Are the attitudes and practices of foodservice managers, catering personnel and students contributing to excessive food wastage at Stellenbosch University? South African Journal of Clinical Nutrition 30(3): 60-67 DOI: 10.1080/16070658.2017.1267348.
- Mashau ME, Moyane JN and Jideani IA. 2012. Assessment of post-harvest losses of fruits at Tshakhuma fruit market in Limpopo Province, South Africa. African Journal of Agricultural Research. Vol. 7(29), pp. 4145-4150. DOI: 10.5897/AJAR12.392.
- Nahman A, and De Lange W. 2013. Cost of food waste along the value chain: evidence form South Africa. *Waste Management*. 33: 2493-2500. DOI: 10.1016/j.wasman.2013.07.012.
- Östergren K, Gustavsson J, Bos-Brouwers H, Timmermans T, Hansen O-J, Møller H, Anderson G, O'Conner C, Soethoudt H, Quested T, Easteal S, Politano A, Bellettato C, Canali M, Falasconi L, Gaiani S, Vittuari M, Schneider F, Moates G, Waldron K and Redingshöfer B. 2014. FUSIONS Definitional Framework for food waste. European Commission (FP7), Coordination and Support Action – CSA. ISBN 987-91-7290-331-9.

NOTES

	• • • • • • • • • • • • • • • • • • • •
•••••••••••••••••••••••••••••••••••••••	
•••••••••••••••••••••••••••••••••••••••	
	 ••••••
	 •
•••••••••••••••••••••••••••••••••••••••	 •
•••••••••••••••••••••••••••••••••••••••	 •

Environment House 473 Steve Biko Road Arcadia Pretoria 0002 Council of Scientific and Industrial Research Meiring Naudé Road Brummeria Pretoria 0184

Call Centre: 086 111 2468

Website: www.environment.gov.za

ISBN: XXX-X-XXX-XXXX-X

Call Centre: 012 841 2911

Website: www.csir.co.za



Find us: Department of Environmental Affairs



Follow us: @EnvironmentZA





To use this QR code conveniently you must have a smartphone equipped with a camera and a QR code reader/ scanner application feature